Self Assessment – Module B Ventilation

1. A patient weighs 180 lbs. Calculate the range of normal tidal volume.

$$180 lbs \times \frac{1 kg}{2.2 lbs} = \frac{180 kg}{2.2} = 81.8 kg$$

- 2. Define Flowrate (\dot{V}). Flowrate is change in volume per unit of time.
- 3. If a patient has a tidal volume of 675 mL and the inspiratory time is 1.6 seconds, calculate the flowrate (\dot{V}) . $\frac{675 mL}{1.6 \, \text{sec}} = 422 \frac{ml}{\text{sec}} + \frac{60 \, \text{sec}}{\text{min}} = 25,320 \frac{ml}{\text{min}}$ $25,320 \frac{ml}{\text{min}} \times \frac{1L}{1000 \, ml} = 25.3 \frac{L}{\text{min}} \text{ (Either answer is OK } \underline{\text{IF}} \text{ you include units.)}$
- 4. Define pressure gradient. A difference in pressure where gas flows from an area of higher pressure to an area of lower pressure.
- 5. The pressure inside the alveoli is called the alveolar (P_A) pressure.
- 6. The difference in pressure between the Airway Pressure (P_{aw}) and the Intrapulmonary pressure (P_{alv}) is called **Transrespiratory Pressure**.
- 7. The difference in pressure between the intrapulmonary pressure and the intrapleural pressure is called **Tran pulmonary Pressure**.
- 8. Which lung pressure is always subatmospheric? Intrapleural
- 9. How is intrapleural pressure measured? **Estimated with an esophageal balloon.**
- 10. Normal breathing is based on which gas law? Boyle's Law
- 11. Explain the 5 steps that occur during normal inspiration
 - A. Inspiratory muscles contract and the diaphragm moves downward.
 - B. Thoracic volume increases.
 - C. Subatmospheric pressure is generated in intrapleural space and alveoli (P_{pl} & P_{alv}).
 - D. A pressure gradient results.
 - E. Gas flow begins (mL/sec) and continues until the pressure is again atmospheric at the end of inspiration ($P_{alv} = P_{bs}$).
- 12. The airway is referred to as a passageway from the larynx down to the alveoli.

| 13. | Explain what is being measured for each of the units listed below | | |
|-----|---|---|-----|
| | A. | ml/cm H ₂ 0: Milliliters of volume per centimeter of water pressure. | |
| | B. | L/cm H ₂ 0 Liters of volume per centimeter of water pressure. | |
| | C. | mL Milliliters of volume. | |
| | D. | L Liters of volume. | |
| | E. | L/min Liters per minute of flow (volume per unit of time). | |
| | F. | mL/sec Milliliters per second of flow (volume per unit of time). | |
| 14. | 500 mL = 0.5 L | | |
| 15. | 8 L = 8,000 mL | | |
| 16. | Com | npliance involves measurement of the | |
| | A. | Elastic properties of the lung B. Inelastic properties of the lungs | |
| 17. | Airway Resistance involves the measurement of the | | |
| | A. | Elastic properties of the lung B. Inelastic properties of the lungs | |
| 18. | What is the normal values for the following | | |
| | A. | Compliance of the lungs? 0.2 L/cm H₂O | |
| | B. | Compliance of the thorax? 0.2 L/cm H₂O | |
| | C. | Total Compliance 0.1 L/cm H ₂ O | |
| 19. | Write | te the formula for compliance Change in Volume divided by Change in Press $\frac{\Delta V}{\Delta P}$ | ure |
| 20. | Based on the compliance formula, if tidal volume stays the same and pressure increases, compliance will | | |
| | A. | Increase B. Decrease C. Stay the same | |
| 21. | An L: | L:S ratio of 2:1 indicates | |
| | A. | Lung maturity B. Lung immaturity C. Transitional | |
| 22. | Surfa | factant is produced from which cells Alveolar Type II. | |
| 23. | When is surfactant usually present in sufficient quantities to support extrauterine life? | | |
| | 35 weeks of gestation | | |

24. Write LaPlace's law.

$$P = \frac{2 \times SurfaceTension}{Radius(r)}$$

- 25. What is the normal total lung compliance? 0.1 L/cm H₂O
- 26. Which of the following L:S ratio's indicate lung immaturity
 - A. 1:2
 - B. 1:1
 - C. 2:1
 - D. 3:1
- 27. If lung compliance decreases than
 - A. Elastance will: a. Increase
- b. Decrease
- c. Stay the same

- B. Pressure will: a. Increase
- b. Decrease
- c. Stay the same

28. Write Poiseuille's Law

$$\overset{\bullet}{\vee} = \frac{\Delta \mathbf{P} \times \mathbf{r}^{4} \times \pi}{\mathbf{8} \times \ell \times \upsilon} \quad \Delta \mathbf{P} = \frac{\vee \times \mathbf{8} \times \ell \times \upsilon}{\mathbf{r}^{4} * \pi}$$

- 29. Patients with emphysema will have
 - A. High compliance
- B. Low compliance
- C. Normal compliance

- 30. Compliance is defined as $\frac{\Delta \mathbf{V}}{\Delta \mathbf{P}}$.
- 31. During which phase of breathing does the alveolar pressure = atmospheric pressure (More than one answer Circle all that apply)
 - A. End of inspiration
 - B. Inspiratory phase
 - C. End of exhalation
 - D. Expiratory phase
- 32. Whose law states that when 1 unit of force or pressure acts upon an elastic body, the elastic body will stretch 1 unit of length or volume? **Hooke's Law**
- 33. Boyles law states that if temperature is held constant, and pressure increases, volume will inversely.
- 34. If the tidal volume is 700 mL and the pressure necessary to push in 700 mL is 25 cm H₂O, the compliance would be 28 mL/cm H₂O or 0.028 L/cm H₂O.

- 35. Name the 4 factors in Poiseuille's Law that have an effect on airway resistance.
 - A. The pressure applied to deliver the breath
 - B. The flow of the gas
 - C. The radius of the tube
 - D. The viscosity of the gas
 - E. The length of the tube
- 36. Name two artificial surfactants that are given to babies born prematurely
 - A. Survanta
 - B. **Exosurf**
 - C. Curosurf
 - D. Infasurf

Pressure to inflate the lungs is 50 cm H_20 Pressure to inflate the lungs is 15 cm H_20

- A. Which one has the lower lung compliance? Patient A
- B. Calculate the lung compliance for patient A and B

I. Patient A:
$$\frac{\Delta V}{\Delta P} = \frac{0.5L}{50cmH_2O} = 0.01 \frac{L}{cmH_2O}$$

II. Patient B:
$$\frac{\Delta V}{\Delta P} = \frac{0.5L}{15cmH_2O} = 0.0\overline{3}\overline{3}\frac{L}{cmH_2O}$$

- 38. The complete absence of spontaneous ventilation is called **APNEA**.
- 39. An increased rate and depth of breathing associated with metabolic disturbances such as diabetes is called **KUSSMAUL'S BREATHING**.
- 40. Define Flow. CHANGE IN VOLUME PER UNIT OF TIME
- 41. Given a V_t of 600 mL and an inspiratory time (T_i) of 0.7 seconds, calculate the flowrate.

$$\dot{V} = \frac{600mL}{0.7\,\text{sec}} = 857\frac{mL}{\text{sec}}$$

- 42. Given a flow (\dot{V}) of 300 mL/sec and an inspiratory time (T_i) of 1.5 seconds, calculate the V_t . $300 \frac{mL}{sec} \times 1.5 \, sec = 450 \, mL$
- 43. Define Compliance and give the normal value for total lung compliance. $\frac{\Delta V}{\Delta P}$, 0.1L/cm H₂O
- 44. Given a V_t of 500 mL and a pressure of 60 cm H₂0, calculate the compliance.

$$\frac{\Delta V}{\Delta P} = \frac{500 \, mL}{60 \, cmH_2 O} = \frac{0.5 \, L}{60 \, cmH_2 O} = 0.008 \, \frac{L}{cmH_2 O}$$

- 45. What is the reciprocal of compliance? **ELASTANCE**
- 46. Does compliance measure the elastic or inelastic properties of the lung? **ELASTIC**
- 47. If pressures increase and Vt stays the same, then compliance will
 - A. Increase

- B. Decrease
- C. Stay the same

- 48. Give examples of high and low compliance.
 - A. HIGH COMPLIANCE WOULD BE FOUND WITH EMPHYSEMA.
 - B. LOW COMPLIANCE WOULD BE FOUND IN PREMATURE BABIES.
- 49. Define Airway Resistance and give the normal value.

THE INELASTIC PROPERTY THAT IMPEDES LUNG EXPANSION. THE NORMAL VALUE IS 0.5 to 2.5 cm H₂O/L/sec

50. Given a transairway pressure of 30 cm H₂0, and a flowrate of 1.2 L/sec, calculate the R_{aw}.

$$R_{aw} = \frac{\Delta P}{\dot{V}} = \frac{30 cm H_2 O}{1.2 \frac{L}{\text{Sec}}} = 25 \frac{cm H_2 O}{L} \frac{L}{\text{Sec}}$$

- 51. What is the reciprocal of resistance? **CONDUCTANCE**
- 52. Does resistance measure the elastic or inelastic properties of the lung? **INELASTIC**
- 53. If flowrate stays the same and the radius of the airway decreases by ½, pressure to breath must (increase or decrease) 16 times?
- 54. If pressure stays the same and the radius of the airway decreases by ½, flow will (increase or decrease) 16 times?
- 55. An increased f is called **TACHYPNEA**.
- 56. Difficulty breathing in a lying down or supine position is called **ORTHOPNEA**.
- 57. How do you assess the presence of hyperventilation and hypoventilation? **ASSESS ARTERIAL BLOOD GASES FOR ABNORMAL CARBON DIOXIDE LEVELS.**
- 58. Normal spontaneous breathing is called **EUPNEA**.